age ~ prev_injuries+injured

24.859 + (−0.089)*prev_injuries + (1.538)*injured

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05, High R and Adj R,
Low AIC

F−statistic: 7.17472 on 2 and 617 Degrees of Freedom


F p−value: 0.00083


R−Squared: 0.02273


Adjusted R−Squared: 0.0195604029209892


Root MSE: 3.82035491265334


AIC: 3429.50950711787



age against prev_injuries
age
prev_injuries
20
25
30
35
0
1
2
3
4



age against injured
age
injured
35
30
25
20
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
363
185
424
Residuals
10
5
0
−5
−10
24.5
25.0
25.5
26.0
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
185
363
424
−3
−2
−1
0
1
2
3


Scale−Location
363
185
424
√|Standardized residuals|
0.0
0.5
1.0
1.5
24.5
25.0
25.5
26.0
Fitted values
lm(formula)


Cook's distance
169
256
381
0.025
0.020
0.015
0.010
0.005
0.000
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
3
2
1
0
−1
−2
−3
381
169
256
Cook's distance
0.000
0.005
0.010
0.015
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
2
169
256
381
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0
0.005
0.01
0.015
0.5
1
1.5
Leverage h_{ii}
lm(formula)