age ~ training_load+injured

24.272 + (0.165)*training_load + (1.422)*injured

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05, High R and Adj R,
Low AIC

F−statistic: 7.56973 on 2 and 617 Degrees of Freedom


F p−value: 0.00057


R−Squared: 0.02395


Adjusted R−Squared: 0.0207856852022125


Root MSE: 3.81796696514048


AIC: 3428.73419152898



age against training_load
age
training_load
20
25
30
35
1
2
3
4
5



age against injured
age
injured
35
30
25
20
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
363
185
424
10
5
0
−5
−10
Residuals
24.5
25.0
25.5
26.0
26.5
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
185
363
424
−3
−2
−1
0
1
2
3


Scale−Location
363
185
424
√|Standardized residuals|
0.0
0.5
1.0
1.5
24.5
25.0
25.5
26.0
26.5
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
156
381
546
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
3
2
1
0
−1
−2
−3
0.000
0.005
0.010
0.015
Leverage
lm(formula)
Cook's distance
546


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
2
546
386
Cook's distance
0.000
0.005
0.010
0.015
0.020
0
0.5
1
1.5
0
0.005
0.01
0.015
Leverage h_{ii}
lm(formula)