bmi ~ age+minutes_played+injured

23.82 + (−0.006)*age + (0)*minutes_played + (−0.106)*injured

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05, High R and Adj R,
Low AIC

F−statistic: 0.34798 on 3 and 616 Degrees of Freedom


F p−value: 0.79063


R−Squared: 0.00169


Adjusted R−Squared: −0.00317004378233232


Root MSE: 1.50578723175117


AIC: 2277.03542155733



bmi against age
bmi
age
20
22
24
26
28
20
25
30
35



bmi against minutes_played
bmi
minutes_played
20
22
24
26
28
0
1000
2000
3000
4000



bmi against injured
bmi
injured
20
22
24
26
28
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
184
4743
Residuals
4
2
0
−2
−4
23.35
23.40
23.45
23.50
23.55
23.60
23.65
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
184
3
474
−3
−2
−1
0
1
2
3


Scale−Location
474
184
√|Standardized residuals|
0.0
0.5
1.0
1.5
23.35
23.40
23.45
23.50
23.55
23.60
23.65
Fitted values
lm(formula)


Cook's distance
43
59
363
0.000
0.005
0.010
0.015
0.020
0.025
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
3
2
1
0
−1
−2
−3
863
43
59
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
2
43
59
363
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0
0.5
1
1.5
0
0.005
0.01
0.015
0.02
0.025
0.03
Leverage h_{ii}
lm(formula)