

bmi ~ age+training_load+prev_injuries


23.876 + (−0.007)*age + (−0.062)*training_load + (0.011)*prev_injuries


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05, High R and Adj R,
Low AIC


F−statistic: 0.4318 on 3 and 616 Degrees of Freedom
F p−value: 0.7303
R−Squared: 0.0021
Adjusted R−Squared: −0.00276138279142502
Root MSE: 1.50548049452941
AIC: 2276.78280093979



bmi against age
bmi
age
20
22
24
26
28
20
25
30
35



bmi against training_load
bmi
training_load
20
22
24
26
28
1
2
3
4
5



bmi against prev_injuries
bmi
prev_injuries
20
22
24
26
28
0
1
2
3
4



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
184
3
474
4
2
0
−2
−4
23.4
23.5
23.6
23.7


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
184
474
3
−3
−2
−1
0
1
2
3


Scale−Location
184
3
474
√|Standardized residuals|
0.0
0.5
1.0
1.5
23.4
23.5
23.6
23.7
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
474
363
271
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
3
2
1
0
−1
−2
−3
271
363
474
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
474
363
271
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0
0.5
1
1.5
2
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)