bmi ~ minutes_played+training_load+prev_injuries

23.782 + (0)*minutes_played + (−0.058)*training_load + (0.011)*prev_injuries

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05, High R and Adj R,
Low AIC

F−statistic: 0.41926 on 3 and 616 Degrees of Freedom


F p−value: 0.73925


R−Squared: 0.00204


Adjusted R−Squared: −0.00282251870014782


Root MSE: 1.50552638656159


AIC: 2276.82059967101



bmi against minutes_played
bmi
minutes_played
20
22
24
26
28
0
1000
2000
3000
4000



bmi against training_load
bmi
training_load
20
22
24
26
28
1
2
3
4
5



bmi against prev_injuries
bmi
prev_injuries
20
22
24
26
28
0
1
2
3
4



Residuals vs Fitted
184
3
474
Residuals
4
2
0
−2
−4
23.4
23.5
23.6
23.7
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
184
474
3
−3
−2
−1
0
1
2
3


Scale−Location
184
3
474
√|Standardized residuals|
0.0
0.5
1.0
1.5
23.4
23.5
23.6
23.7
Fitted values
lm(formula)


Cook's distance
43
220
474
0.000
0.005
0.010
0.015
0.020
0.025
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
220
43
474
3
2
1
0
−1
−2
−3
0.000
0.005
0.010
0.015
0.020
0.025


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
474
220
43
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0
0.5
1
1.5
2
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)