


bmi ~ minutes_played+training_load
23.79 + (0)*minutes_played + (−0.058)*training_load


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 0.61209 on 2 and 617 Degrees of Freedom
F p−value: 0.54255
R−Squared: 0.00198
Adjusted R−Squared: −0.0012549317016457
Root MSE: 1.50556979334314
AIC: 2274.85635037834



bmi against minutes_played
bmi
minutes_played
20
22
24
26
28
0
1000
2000
3000
4000



bmi against training_load
bmi
training_load
20
22
24
26
28
1
2
3
4
5



Residuals vs Fitted
184
3
474
Residuals
4
2
0
−2
−4
23.40
23.45
23.50
23.55
23.60
23.65
23.70
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
184
474
3
−3
−2
−1
0
1
2
3


Scale−Location
184
3
474
√|Standardized residuals|
0.0
0.5
1.0
1.5
23.40
23.45
23.50
23.55
23.60
23.65
23.70
Fitted values
lm(formula)


Cook's distance
43
474
220
0.035
0.030
0.025
0.020
0.015
0.010
0.005
0.000
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
220
43
474
3
2
1
0
−1
−2
−3
0.000
0.005
0.010
0.015
0.020
0.025


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
474
43
220
Cook's distance
Leverage h_{ii}
lm(formula)
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0.035
0
0.005
0.01
0.015
0.02
0.025
0
0.5
1
1.5
2