bmi ~ prev_injuries

23.514 + (0.007)*prev_injuries

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC

F−statistic: 0.01666 on 1 and 618 Degrees of Freedom


F p−value: 0.89733


R−Squared: 3e−05


Adjusted R−Squared: −0.00159111743669205


Root MSE: 1.50704231401703


AIC: 2274.06853815158



bmi against prev_injuries
bmi
prev_injuries
20
22
24
26
28
0
1
2
3
4



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
184
474
3
4
2
0
−2
−4
23.515
23.520
23.525
23.530
23.535
23.540


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
3
474
184
−3
−2
−1
0
1
2
3


Scale−Location
474
184
3
√|Standardized residuals|
0.0
0.5
1.0
1.5
23.515
23.520
23.525
23.530
23.535
23.540
Fitted values
lm(formula)


Cook's distance
271
220
534
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
3
2
1
0
−1
−2
−3
220
271
534
Cook's distance
0.000
0.005
0.010
0.015
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
271
534
220
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0
0.002
0.004
0.006
0.008
0.01
0.012
0.014
Leverage h_{ii}
lm(formula)
2
1.5
1
0.5
0