bmi ~ training_load+injured

23.723 + (−0.058)*training_load + (−0.098)*injured

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05, High R and Adj R,
Low AIC

F−statistic: 0.73562 on 2 and 617 Degrees of Freedom


F p−value: 0.47963


R−Squared: 0.00238


Adjusted R−Squared: −0.000854958083650104


Root MSE: 1.50526904658539


AIC: 2274.60862806527



bmi against training_load
bmi
training_load
20
22
24
26
28
1
2
3
4
5



bmi against injured
bmi
injured
20
22
24
26
28
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
184
3
474
4
2
0
−2
−4
23.35
23.40
23.45
23.50
23.55
23.60
23.65


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
184
474
3
−3
−2
−1
0
1
2
3


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
184
3
474
23.35
23.40
23.45
23.50
23.55
23.60
23.65
Fitted values
lm(formula)


Cook's distance
474
101
59
0.030
0.025
0.020
0.015
0.010
0.005
0.000
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
3
2
1
0
−1
−2
−3
0.000
0.005
0.010
0.015
Leverage
lm(formula)
Cook's distance
59
474
101


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
474
101
59
Cook's distance
Leverage h_{ii}
lm(formula)
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0
0.005
0.01
0.015
0
0.5
1
1.5
2