


bmi ~ training_load+prev_injuries
23.714 + (−0.063)*training_load + (0.011)*prev_injuries


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 0.55836 on 2 and 617 Degrees of Freedom
F p−value: 0.57243
R−Squared: 0.00181
Adjusted R−Squared: −0.00142897703191136
Root MSE: 1.50570064213964
AIC: 2274.96411386969



bmi against training_load
bmi
training_load
20
22
24
26
28
1
2
3
4
5



bmi against prev_injuries
bmi
prev_injuries
20
22
24
26
28
0
1
2
3
4



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
184
3
474
4
2
0
−2
−4
23.40
23.45
23.50
23.55
23.60
23.65


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
184
474
3
−3
−2
−1
0
1
2
3


Scale−Location
184
3
474
√|Standardized residuals|
0.0
0.5
1.0
1.5
23.40
23.45
23.50
23.55
23.60
23.65
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0.035
474
271
507
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
507
271
474
3
2
1
0
−1
−2
−3
0.000
0.005
0.010
0.015


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
474
507
271
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0
0.5
1
1.5
2
0
0.005
0.01
0.015
Leverage h_{ii}
lm(formula)