bmi ~ training_load

23.722 + (−0.063)*training_load

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05, High R and Adj R,
Low AIC

F−statistic: 1.08068 on 1 and 618 Degrees of Freedom


F p−value: 0.29895


R−Squared: 0.00175


Adjusted R−Squared: 0.000130314283576571


Root MSE: 1.50574668247231


AIC: 2273.00202920166



bmi against training_load
bmi
training_load
20
22
24
26
28
1
2
3
4
5



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
184
3
474
4
2
0
−2
−4
23.40
23.45
23.50
23.55
23.60
23.65


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
184
474
3
−3
−2
−1
0
1
2
3


Scale−Location
184
3
474
√|Standardized residuals|
0.0
0.5
1.0
1.5
23.40
23.45
23.50
23.55
23.60
23.65
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.00
0.01
0.02
0.03
0.04
0.05
474
507
101
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
3
2
1
0
−1
−2
−3
507
101
474
Cook's distance
0.000
0.002
0.004
0.006
0.008
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
Cook's distance
Leverage h_{ii}
lm(formula)
0.00
0.01
0.02
0.03
0.04
0
0.002
0.004
0.006
0.008
0
0.5
1
1.5
2
2.5
3
3.5
474
507
101