goals_conceded ~ goals_scored+opponent_rank
1.37 + (−0.026)*goals_scored + (−0.04)*opponent_rank

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC

F−statistic: 3.58275 on 2 and 43 Degrees of Freedom
F p−value: 0.03638
R−Squared: 0.14284
Adjusted R−Squared: 0.102969005896273
Root MSE: 0.730132715134113
AIC: 109.605680752308


goals_conceded against goals_scored
goals_conceded
goals_scored
0.0
0.5
1.0
1.5
2.0
2.5
3.0
0
2
4
6
8


goals_conceded against opponent_rank
goals_conceded
0.0
0.5
1.0
1.5
2.0
2.5
3.0
5
10
15
20
opponent_rank


Residuals vs Fitted
Residuals
Fitted values
lm(formula)
34
29
44
2.0
1.5
1.0
0.5
0.0
−0.5
−1.0
−1.5
0.2
0.4
0.6
0.8
1.0
1.2


Q−Q Residuals
Standardized residuals
3
2
1
0
−1
−2
−1
0
1
2
34
29
44
Theoretical Quantiles
lm(formula)


Scale−Location
34
29
44
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.2
0.4
0.6
0.8
1.0
1.2
Fitted values
lm(formula)


Cook's distance
29
34
44
0.00
0.05
0.10
0.15
Cook's distance
0
10
20
30
40
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
34
29
44
1
0.5
0.5
1
3
2
1
0
−1
−2
0.0
0.1
0.2
0.3
0.4
0.5


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
2
1.5
1
34
29
44
Cook's distance
0.00
0.05
0.10
0.15
0.5
0
0
0.1
0.2
0.3
0.4
0.5
Leverage h_{ii}
lm(formula)