goals_conceded ~ goals_scored
0.951 + (−0.095)*goals_scored

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC

F−statistic: 1.29432 on 1 and 44 Degrees of Freedom
F p−value: 0.26141
R−Squared: 0.02858
Adjusted R−Squared: 0.00649787418791004
Root MSE: 0.77727481249699
AIC: 113.361904800836


goals_conceded against goals_scored
goals_conceded
goals_scored
0.0
0.5
1.0
1.5
2.0
2.5
3.0
0
2
4
6
8


Residuals vs Fitted
34
29
42
Residuals
2
1
0
−1
0.2
0.4
0.6
0.8
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
34
29
42
−2
−1
0
1
2
3


Scale−Location
34
29
42
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.2
0.4
0.6
0.8
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.00
0.02
0.04
0.06
0.08
0.10
0.12
8
29
34
0
10
20
30
40
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
34
29
8
1
0.5
0.5
1
3
2
1
0
−1
0.0
0.1
0.2
0.3
0.4
0.5


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
Cook's distance
Leverage h_{ii}
lm(formula)
0.00
0.02
0.04
0.06
0.08
0.10
0.12
0
0.1
0.2
0.3
0.4
0.5
3
2.5
2
1.5
1
0.5
0
34
29
8