goals_conceded ~ opponent_rank
1.359 + (−0.042)*opponent_rank

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC

F−statistic: 7.22202 on 1 and 44 Degrees of Freedom
F p−value: 0.01013
R−Squared: 0.14099
Adjusted R−Squared: 0.121471516433526
Root MSE: 0.730917091975323
AIC: 107.70446270481


goals_conceded against opponent_rank
goals_conceded
0.0
0.5
1.0
1.5
2.0
2.5
3.0
5
10
15
20
opponent_rank


Residuals vs Fitted
Residuals
Fitted values
lm(formula)
44
34
29
2.0
1.5
1.0
0.5
0.0
−0.5
−1.0
−1.5
0.4
0.6
0.8
1.0
1.2


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
34
29
44
3
2
1
0
−1
−2
−1
0
1
2


Scale−Location
34
29
44
1.5
1.0
0.5
0.0
√|Standardized residuals|
0.4
0.6
0.8
1.0
1.2
Fitted values
lm(formula)


Cook's distance
29
34
44
0.00
0.05
0.10
0.15
0.20
Cook's distance
0
10
20
30
40
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
34
29
44
3
2
1
0
−1
−2
0.00
0.02
0.04
0.06
0.08


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
29
34
44
Cook's distance
0.00
0.05
0.10
0.15
0.20
0.02
0.03
0.04
0.05
0.06
0.07
0.08
Leverage h_{ii}
lm(formula)
0
0.5
1
1.5
2