goals_scored ~ goals_conceded+opponent_rank
0.556 + (−0.083)*goals_conceded + (0.065)*opponent_rank

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC

F−statistic: 2.91527 on 2 and 43 Degrees of Freedom
F p−value: 0.06497
R−Squared: 0.1194
Adjusted R−Squared: 0.078445619402766
Root MSE: 1.31069145317207
AIC: 163.433388937384


goals_scored against goals_conceded
goals_scored
goals_conceded
0
2
4
6
8
0.0
0.5
1.0
1.5
2.0
2.5
3.0


goals_scored against opponent_rank
goals_scored
0
2
4
6
8
5
10
15
20
opponent_rank


Residuals vs Fitted
Residuals
Fitted values
lm(formula)
8
19
46
6
4
2
0
−2
0.5
1.0
1.5
2.0


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
8
19
46
−2
−1
0
1
2
−1
0
1
2
3
4
5


Scale−Location
√|Standardized residuals|
Fitted values
lm(formula)
0.0
0.5
1.0
1.5
2.0
0.5
1.0
1.5
2.0
8
19
46


Cook's distance
8
34
44
Cook's distance
0.0
0.1
0.2
0.3
0.4
0.5
0.6
0
10
20
30
40
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
1
0.5
8
34
44
5
4
3
2
1
0
−1
−2
0.00
0.05
0.10
0.15
0.20


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
5
4
3
8
34
44
Cook's distance
0.0
0.1
0.2
0.3
0.4
0.5
0.6
0
1
2
0
0.05
0.1
0.15
Leverage h_{ii}
lm(formula)