goals_scored ~ goals_conceded

1.552 + (−0.299)*goals_conceded

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05, High R and Adj R,
Low AIC

F−statistic: 1.29432 on 1 and 44 Degrees of Freedom
F p−value: 0.26141
R−Squared: 0.02858
Adjusted R−Squared: 0.00649787418791004
Root MSE: 1.37662766931489
AIC: 165.948929783538


goals_scored against goals_conceded
goals_scored
goals_conceded
0
2
4
6
8
0.0
0.5
1.0
1.5
2.0
2.5
3.0


Residuals vs Fitted
Residuals
6
4
2
0
−2
8
19
9
0.8
1.0
1.2
1.4
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
8
5
19
−2
−1
0
1
2
−1
0
1
2
3
4
5


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
2.0
0.8
1.0
1.2
1.4
Fitted values
lm(formula)
8
19


Cook's distance
8
7
34
0.0
0.1
0.2
0.3
0.4
0.5
Cook's distance
0
10
20
30
40
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
5
4
3
2
1
0
−1
8
7
34
1
0.5
Cook's distance
0.00
0.05
0.10
0.15
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
5
4
3
8
7
34
Cook's distance
0.0
0.1
0.2
0.3
0.4
0.5
0
0.05
0.1
0.15
Leverage h_{ii}
lm(formula)
0
1
2