goals_scored ~ opponent_rank
0.443 + ( 0.068)*opponent_rank

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC

F−statistic: 5.85895 on 1 and 44 Degrees of Freedom
F p−value: 0.01969
R−Squared: 0.11751
Adjusted R−Squared: 0.0974539588578466
Root MSE: 1.31209952050083
AIC: 161.532170889886


goals_scored against opponent_rank
goals_scored
0
2
4
6
8
5
10
15
20
opponent_rank


Residuals vs Fitted
Residuals
Fitted values
lm(formula)
6
4
2
0
−2
0.5
1.0
1.5
2.0
8
19
46


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
8
19
46
−2
−1
0
1
2
−1
0
1
2
3
4
5


Scale−Location
√|Standardized residuals|
Fitted values
lm(formula)
0.0
0.5
1.0
1.5
2.0
0.5
1.0
1.5
2.0
8
19
46


Cook's distance
Cook's distance
1.0
0.8
0.6
0.4
0.2
0.0
8
1
46
0
10
20
30
40
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
1
0.5
8
1
46
5
4
3
2
1
0
−1
−2
0.00
0.02
0.04
0.06
0.08


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
Cook's distance
Leverage $h_{ii}$
lm(formula)
0.0
0.2
0.4
0.6
0.8
0.02
0.03
0.04
0.05
0.06
0.07
0.08
0
1
2
3
4
5
8
46
1