injured ~ age+minutes_played+bmi

−0.092 + (0.015)*age + (0)*minutes_played + (−0.007)*bmi

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05, High R and Adj R,
Low AIC

F−statistic: 5.25242 on 3 and 616 Degrees of Freedom


F p−value: 0.00139


R−Squared: 0.02494


Adjusted R−Squared: 0.0201932948153286


Root MSE: 0.383811651739812


AIC: 582.055643312376



injured against age
injured
0.0
0.2
0.4
0.6
0.8
1.0
20
25
30
35
age



injured against minutes_played
injured
0.0
0.2
0.4
0.6
0.8
1.0
0
1000
2000
3000
4000
minutes_played



injured against bmi
injured
bmi
0.0
0.2
0.4
0.6
0.8
1.0
20
22
24
26
28



Residuals vs Fitted
546
361
169
1.0
0.5
0.0
−0.5
Residuals
0.0
0.1
0.2
0.3
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
546
169
361
3
2
1
0
−1
−3
−2
−1
0
1
2
3


Scale−Location
546
361
169
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.0
0.1
0.2
0.3
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.00
0.01
0.02
0.03
0.04
169
447
546
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
546
447
169
Cook's distance
3
2
1
0
−1
0.000
0.005
0.010
0.015
0.020
0.025
0.030


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
2
1.5
1
0.5
0
546
169
447
Cook's distance
0.00
0.01
0.02
0.03
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)