

injured ~ age+minutes_played+training_load


−0.35 + (0.014)*age + (0)*minutes_played + (0.047)*training_load


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05, High R and Adj R,
Low AIC


F−statistic: 8.22464 on 3 and 616 Degrees of Freedom
F p−value: 2e−05
R−Squared: 0.03851
Adjusted R−Squared: 0.033829893145519
Root MSE: 0.381131416967338
AIC: 573.366095411073



injured against age
injured
0.0
0.2
0.4
0.6
0.8
1.0
20
25
30
35
age



injured against minutes_played
injured
0.0
0.2
0.4
0.6
0.8
1.0
0
1000
2000
3000
4000
minutes_played



injured against training_load
injured
training_load
0.0
0.2
0.4
0.6
0.8
1.0
1
2
3
4
5



Residuals vs Fitted
546
238
281
1.0
0.5
0.0
−0.5
Residuals
0.0
0.1
0.2
0.3
0.4
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
546
238
281
3
2
1
0
−1
−3
−2
−1
0
1
2
3


Scale−Location
546
238
281
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.0
0.1
0.2
0.3
0.4
Fitted values
lm(formula)


Cook's distance
546
281
480
0.00
0.01
0.02
0.03
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
3
2
1
0
−1
546
281
480
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
546
281
480
2
1.5
1
0.5
0
Cook's distance
0.035
0.030
0.025
0.020
0.015
0.010
0.005
0.000
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)