


injured ~ age+minutes_played
−0.255 + (0.015)*age + (0)*minutes_played


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 7.66223 on 2 and 617 Degrees of Freedom
F p−value: 0.00052
R−Squared: 0.02424
Adjusted R−Squared: 0.0210722025338561
Root MSE: 0.38395073885672
AIC: 580.504917798767



injured against age
injured
0.0
0.2
0.4
0.6
0.8
1.0
20
25
30
35
age



injured against minutes_played
injured
0.0
0.2
0.4
0.6
0.8
1.0
0
1000
2000
3000
4000
minutes_played



Residuals vs Fitted
546
361
72
1.0
0.5
0.0
−0.5
Residuals
0.05
0.10
0.15
0.20
0.25
0.30
0.35
Fitted values
lm(formula)


Q−Q Residuals
546
361
72
Standardized residuals
−1
0
1
2
−3
−2
−1
0
1
2
3
Theoretical Quantiles
lm(formula)


Scale−Location
546
361
72
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.05
0.10
0.15
0.20
0.25
0.30
0.35
Fitted values
lm(formula)


Cook's distance
546
169
480
0.00
0.01
0.02
0.03
0.04
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
546
169
480
Cook's distance
Standardized residuals
Leverage
lm(formula)
3
2
1
0
−1
0.000
0.005
0.010
0.015
0.020


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
546
169
480
Cook's distance
0.00
0.01
0.02
0.03
0.04
2
1.5
1
0.5
0
0
0.005
0.01
0.015
0.02
Leverage h_{ii}
lm(formula)