


injured ~ age+training_load+prev_injuries
−0.379 + (0.014)*age + (0.044)*training_load + (0.082)*prev_injuries


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 20.42653 on 3 and 616 Degrees of Freedom
F p−value: 0
R−Squared: 0.09048
Adjusted R−Squared: 0.0860495384847297
Root MSE: 0.370688642187018
AIC: 538.916699103353



injured against age
injured
0.0
0.2
0.4
0.6
0.8
1.0
20
25
30
35
age



injured against training_load
injured
training_load
0.0
0.2
0.4
0.6
0.8
1.0
1
2
3
4
5



injured against prev_injuries
injured
0.0
0.2
0.4
0.6
0.8
1.0
0
1
2
3
4
prev_injuries



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
1.0
0.5
0.0
−0.5
0.0
0.1
0.2
0.3
0.4
0.5
288
101


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
238
281
101
3
2
1
0
−1
−3
−2
−1
0
1
2
3


Scale−Location
√|Standardized residuals|
Fitted values
lm(formula)
0.0
0.5
1.0
1.5
0.0
0.1
0.2
0.3
0.4
0.5
288
101


Cook's distance
169
238
281
0.000
0.005
0.010
0.015
0.020
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
169
0.000
0.005
0.010
0.015
0.020
0.025
−2
−1
0
1
2
3


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
2
288
169
0.020
0.015
0.010
0.005
0.000
Cook's distance
1.5
1
0.5
0
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)