


injured ~ age+training_load
−0.327 + (0.014)*age + (0.049)*training_load


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 12.23527 on 2 and 617 Degrees of Freedom
F p−value: 1e−05
R−Squared: 0.03815
Adjusted R−Squared: 0.0350297440528027
Root MSE: 0.381203729338722
AIC: 571.601339296178



injured against age
injured
0.0
0.2
0.4
0.6
0.8
1.0
20
25
30
35
age



injured against training_load
injured
training_load
0.0
0.2
0.4
0.6
0.8
1.0
1
2
3
4
5



Residuals vs Fitted
546
288
1.0
0.5
0.0
−0.5
Residuals
0.0
0.1
0.2
0.3
0.4
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
546
238
281
3
2
1
0
−1
−3
−2
−1
0
1
2
3


Scale−Location
546
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.0
0.1
0.2
0.3
0.4
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
238
281
546
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
546
0.000
0.005
0.010
0.015
3
2
1
0
−1


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
546
288
0.025
0.020
0.015
0.010
0.005
0.000
Cook's distance
2
1.5
1
0.5
0
0
0.005
0.01
0.015
Leverage h_{ii}
lm(formula)