


injured ~ minutes_played+bmi
0.306 + (0)*minutes_played + (−0.008)*bmi


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 0.85022 on 2 and 617 Degrees of Freedom
F p−value: 0.42782
R−Squared: 0.00275
Adjusted R−Squared: −0.000484161404904038
Root MSE: 0.388155090480626
AIC: 594.009407324778



injured against minutes_played
injured
0.0
0.2
0.4
0.6
0.8
1.0
0
1000
2000
3000
4000
minutes_played



injured against bmi
injured
bmi
0.0
0.2
0.4
0.6
0.8
1.0
20
22
24
26
28



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
1.0
0.8
0.6
0.4
0.2
0.0
−0.2
0.10
0.12
0.14
0.16
0.18
0.20
0.22
0.24
546
480
238


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
2.5
2.0
1.5
1.0
0.5
0.0
−0.5
−3
−2
−1
0
1
2
3
480
546
238


Scale−Location
546
480
238
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.10
0.12
0.14
0.16
0.18
0.20
0.22
0.24
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
59
480
546
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
2.5
2.0
1.5
1.0
0.5
0.0
−0.5
59
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
2.5
2
480
546
59
1.5
1
0.5
0
Cook's distance
0.025
0.020
0.015
0.010
0.005
0.000
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)