injured ~ minutes_played+prev_injuries+bmi

0.236 + (0)*minutes_played + (0.085)*prev_injuries + (−0.008)*bmi

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05, High R and Adj R,
Low AIC

F−statistic: 12.97618 on 3 and 616 Degrees of Freedom


F p−value: 0


R−Squared: 0.05944


Adjusted R−Squared: 0.0548586986885331


Root MSE: 0.37696091429089


AIC: 559.722703236846



injured against minutes_played
injured
0.0
0.2
0.4
0.6
0.8
1.0
0
1000
2000
3000
4000
minutes_played



injured against prev_injuries
injured
0.0
0.2
0.4
0.6
0.8
1.0
0
1
2
3
4
prev_injuries



injured against bmi
injured
bmi
0.0
0.2
0.4
0.6
0.8
1.0
20
22
24
26
28



Residuals vs Fitted
Residuals
1.0
0.5
0.0
−0.5
0.1
0.2
0.3
0.4
0.5
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
−1
0
1
2
238


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.1
0.2
0.3
0.4
0.5
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.000
0.005
0.010
0.015
0.020
169
480
546
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
169
546
480
2
1
0
−1
0.000
0.005
0.010
0.015
0.020
0.025


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
2.5
2
546
480
169
Cook's distance
0.000
0.005
0.010
0.015
1.5
1
0.5
0
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)