


injured ~ minutes_played+prev_injuries
0.048 + (0)*minutes_played + (0.085)*prev_injuries


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 19.1654 on 2 and 617 Degrees of Freedom
F p−value: 0
R−Squared: 0.05849
Adjusted R−Squared: 0.0554388607983113
Root MSE: 0.377150957175498
AIC: 558.347685260919



injured against minutes_played
injured
0.0
0.2
0.4
0.6
0.8
1.0
0
1000
2000
3000
4000
minutes_played



injured against prev_injuries
injured
0.0
0.2
0.4
0.6
0.8
1.0
0
1
2
3
4
prev_injuries



Residuals vs Fitted
Residuals
1.0
0.5
0.0
−0.5
0.1
0.2
0.3
0.4
0.5
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
−1
0
1
2
72


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.1
0.2
0.3
0.4
0.5
Fitted values
lm(formula)


Cook's distance
480
546
97
Cook's distance
0.000
0.005
0.010
0.015
0.020
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
546
480
97
Standardized residuals
2
1
0
−1
Cook's distance
0.000
0.005
0.010
0.015
0.020
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
2.5
2
480
546
97
0.020
0.015
0.010
0.005
0.000
Cook's distance
1.5
1
0.5
0
0
0.005
0.01
0.015
0.02
Leverage h_{ii}
lm(formula)