


injured ~ minutes_played+training_load
0.001 + ( 0)*minutes_played + ( 0.051)*training_load


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 5.84329 on 2 and 617 Degrees of Freedom
F p−value: 0.00306
R−Squared: 0.01859
Adjusted R−Squared: 0.0154076584462408
Root MSE: 0.385059997732716
AIC: 584.082194148136



injured against minutes_played
injured
0.0
0.2
0.4
0.6
0.8
1.0
0
1000
2000
3000
4000
minutes_played



injured against training_load
injured
training_load
0.0
0.2
0.4
0.6
0.8
1.0
1
2
3
4
5



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
1.0
0.8
0.6
0.4
0.2
0.0
−0.2
−0.4
0.05
0.10
0.15
0.20
0.25
281


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
2816
1101
238
2.5
2.0
1.5
1.0
0.5
0.0
−0.5
−3
−2
−1
0
1
2
3


Scale−Location
√|Standardized residuals|
Fitted values
lm(formula)
0.0
0.5
1.0
1.5
0.05
0.10
0.15
0.20
0.25
281


Cook's distance
480
281
546
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
281
546
480
Cook's distance
Standardized residuals
Leverage
lm(formula)
2
1
0
−1
0.000
0.005
0.010
0.015
0.020
0.025


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
2.5
2
480
281
546
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0
0.5
1
1.5
0
0.005
0.01
0.015
0.02
0.025
Leverage $h_{ii}$
lm(formula)