


injured ~ minutes_played
0.125 + (0)*minutes_played


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 1.16032 on 1 and 618 Degrees of Freedom
F p−value: 0.28182
R−Squared: 0.00187
Adjusted R−Squared: 0.000258934920428278
Root MSE: 0.38832522136471
AIC: 592.552788261606



injured against minutes_played
injured
0.0
0.2
0.4
0.6
0.8
1.0
0
1000
2000
3000
4000
minutes_played



Residuals vs Fitted
480
546
97
Residuals
0.8
0.6
0.4
0.2
0.0
−0.2
0.14
0.16
0.18
0.20
0.22
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
2.5
2.0
1.5
1.0
0.5
0.0
−0.5
480
546
−3
−2
−1
0
1
2
3
Theoretical Quantiles
lm(formula)


Scale−Location
480
546
97
1.5
1.0
0.5
0.0
√|Standardized residuals|
0.14
0.16
0.18
0.20
0.22
Fitted values
lm(formula)


Cook's distance
480
546
97
0.00
0.01
0.02
0.03
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
2.5
2.0
1.5
1.0
0.5
0.0
−0.5
97
546
480
Cook's distance
0.000
0.005
0.010
0.015
0.020
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
2.5
2
480
546
97
1.5
1
0.5
0
Cook's distance
0.00
0.01
0.02
0.03
0
0.005
0.01
0.015
0.02
Leverage h_{ii}
lm(formula)