


injured ~ prev_injuries+bmi
0.301 + (0.085)*prev_injuries + (−0.008)*bmi


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 18.85701 on 2 and 617 Degrees of Freedom
F p−value: 0
R−Squared: 0.0576
Adjusted R−Squared: 0.054549036953835
Root MSE: 0.377328562891231
AIC: 558.931481336807



injured against prev_injuries
injured
0.0
0.2
0.4
0.6
0.8
1.0
0
1
2
3
4
prev_injuries



injured against bmi
injured
bmi
0.0
0.2
0.4
0.6
0.8
1.0
20
22
24
26
28



Residuals vs Fitted
Residuals
1.0
0.5
0.0
−0.5
0.1
0.2
0.3
0.4
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
169
238
−3
−2
−1
0
1
2
3
−1
0
1
2


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.1
0.2
0.3
0.4
Fitted values
lm(formula)


Cook's distance
59
101
271
0.000
0.005
0.010
0.015
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
101
59
271
Cook's distance
0.000
0.005
0.010
0.015
0.020
−1
0
1
2


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
2.5
2
101
59
271
Cook's distance
0.000
0.005
0.010
0.015
0
0.5
1
1.5
0
0.005
0.01
0.015
0.02
Leverage h_{ii}
lm(formula)