


injured ~ prev_injuries
0.108 + (0.085)*prev_injuries


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 37.0709 on 1 and 618 Degrees of Freedom
F p−value: 0
R−Squared: 0.05659
Adjusted R−Squared: 0.0550641158986734
Root MSE: 0.377531334860741
AIC: 557.597663893303



injured against prev_injuries
injured
0.0
0.2
0.4
0.6
0.8
1.0
0
1
2
3
4
prev_injuries



Residuals vs Fitted
Residuals
1.0
0.5
0.0
−0.5
0.10
0.15
0.20
0.25
0.30
0.35
0.40
0.45
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
−1
0
1
2


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.10
0.15
0.20
0.25
0.30
0.35
0.40
0.45
Fitted values
lm(formula)


Cook's distance
34
97
181
Cook's distance
0.000
0.005
0.010
0.015
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
2
1
0
−1
Cook's distance
0.000
0.005
0.010
0.015
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
2.5
2
1.5
0.5
1
0
Cook's distance
0.000
0.005
0.010
0.015
0
0.002
0.004
0.006
0.008
0.01
0.012
0.014
Leverage h_{ii}
lm(formula)