


injured ~ training_load+bmi
0.17 + (0.052)*training_load + (−0.006)*bmi


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 5.97255 on 2 and 617 Degrees of Freedom
F p−value: 0.0027
R−Squared: 0.01899
Adjusted R−Squared: 0.0158123511991992
Root MSE: 0.384980854822556
AIC: 583.82730582989



injured against training_load
injured
training_load
0.0
0.2
0.4
0.6
0.8
1.0
1
2
3
4
5



injured against bmi
injured
bmi
0.0
0.2
0.4
0.6
0.8
1.0
20
22
24
26
28



Residuals vs Fitted
238
281
101
1.0
0.5
0.0
Residuals
0.05
0.10
0.15
0.20
0.25
0.30
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
2.5
2.0
1.5
1.0
0.5
0.0
−0.5
−3
−2
−1
0
1
2
3
238
281
101


Scale−Location
238
281
101
1.5
1.0
0.5
0.0
√|Standardized residuals|
0.05
0.10
0.15
0.20
0.25
0.30
Fitted values
lm(formula)


Cook's distance
101
238
281
0.035
0.030
0.025
0.020
0.015
0.010
0.005
0.000
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
281
238
101
Standardized residuals
2
1
0
−1
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
2.5
2
101
238
281
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0
0.5
1
1.5
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)