


injured ~ training_load+prev_injuries+bmi
0.125 + (0.046)*training_load + (0.083)*prev_injuries + (−0.007)*bmi


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 15.91474 on 3 and 616 Degrees of Freedom
F p−value: 0
R−Squared: 0.07193
Adjusted R−Squared: 0.0674118245747137
Root MSE: 0.374449197020003
AIC: 551.432846688173



injured against training_load
injured
training_load
0.0
0.2
0.4
0.6
0.8
1.0
1
2
3
4
5



injured against prev_injuries
injured
0.0
0.2
0.4
0.6
0.8
1.0
0
1
2
3
4
prev_injuries



injured against bmi
injured
bmi
0.0
0.2
0.4
0.6
0.8
1.0
20
22
24
26
28



Residuals vs Fitted
238
281
101
1.0
0.5
0.0
−0.5
Residuals
0.0
0.1
0.2
0.3
0.4
0.5
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
238
281
101
3
2
1
0
−1
−3
−2
−1
0
1
2
3


Scale−Location
238
281
101
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.0
0.1
0.2
0.3
0.4
0.5
Fitted values
lm(formula)


Cook's distance
101
238
281
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
3
2
1
0
−1
281
238
101
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
2
1.5
1
0.5
0
101
238
281
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)