


injured ~ training_load+prev_injuries
−0.039 + (0.047)*training_load + (0.083)*prev_injuries


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 23.65472 on 2 and 617 Degrees of Freedom
F p−value: 0
R−Squared: 0.07122
Adjusted R−Squared: 0.0682053370134692
Root MSE: 0.374593542601808
AIC: 549.910759400457



injured against training_load
injured
training_load
0.0
0.2
0.4
0.6
0.8
1.0
1
2
3
4
5



injured against prev_injuries
injured
0.0
0.2
0.4
0.6
0.8
1.0
0
1
2
3
4
prev_injuries



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
1.0
0.5
0.0
−0.5
0.0
0.1
0.2
0.3
0.4
0.5


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
3
2
1
0
−1
−3
−2
−1
0
1
2
3
1012
38
281


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.0
0.1
0.2
0.3
0.4
0.5
Fitted values
lm(formula)


Cook's distance
101
238
281
0.025
0.020
0.015
0.010
0.005
0.000
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
3
2
1
0
−1
0.000
0.005
0.010
0.015
Leverage
lm(formula)
Cook's distance


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
Cook's distance
Leverage h_{ii}
lm(formula)
0.000
0.005
0.010
0.015
0.020
0.025
0
0.005
0.01
0.015
3
2.5
2
1.5
1
0.5
0