


injured ~ training_load
0.018 + (0.052)*training_load


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 11.56486 on 1 and 618 Degrees of Freedom
F p−value: 0.00072
R−Squared: 0.01837
Adjusted R−Squared: 0.0167812157551445
Root MSE: 0.385103012833084
AIC: 582.220706966279



injured against training_load
injured
training_load
0.0
0.2
0.4
0.6
0.8
1.0
1
2
3
4
5



Residuals vs Fitted
Residuals
1.0
0.8
0.6
0.4
0.2
0.0
−0.2
−0.4
0.10
0.15
0.20
0.25
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
2.5
2.0
1.5
1.0
0.5
0.0
−0.5
−3
−2
−1
0
1
2
3
281


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.10
0.15
0.20
0.25
Fitted values
lm(formula)


Cook's distance
101
238
281
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
2
1
0
−1
Cook's distance
0.000
0.002
0.004
0.006
0.008
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
Cook's distance
Leverage h_{ii}
lm(formula)
0.000
0.005
0.010
0.015
0.020
0.025
0
0.002
0.004
0.006
0.008
2.5
2
1.5
1
0.5
0