


minutes_played ~ age+bmi
2885.971 + (−3.11)*age + (−11.383)*bmi


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 0.26421 on 2 and 617 Degrees of Freedom
F p−value: 0.7679
R−Squared: 0.00086
Adjusted R−Squared: −0.00238301871038127
Root MSE: 709.161815444123
AIC: 9906.94760763758



minutes_played against age
minutes_played
0
1000
2000
3000
4000
20
25
30
35
age



minutes_played against bmi
minutes_played
0
1000
2000
3000
4000
20
22
24
26
28
bmi



Residuals vs Fitted
Residuals
1000
0
−1000
−2000
43
480
466
2480
2500
2520
2540
2560
2580
2600
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
43
480
466


Scale−Location
43
480
466
√|Standardized residuals|
0.0
0.5
1.0
1.5
2480
2500
2520
2540
2560
2580
2600
Fitted values
lm(formula)


Cook's distance
43
220
546
0.00
0.01
0.02
0.03
0.04
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
220
546
43
2
1
0
−1
−2
−3
−4
0.000
0.005
0.010
0.015
0.020


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
43
546
220
Cook's distance
0.00
0.01
0.02
0.03
0.04
0
0.005
0.01
0.015
0.02
Leverage h_{ii}
lm(formula)
2
1.5
1
0.5
0