


minutes_played ~ age+injured
2632.24 + (−4.301)*age + (85.38)*injured


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 0.74553 on 2 and 617 Degrees of Freedom
F p−value: 0.47491
R−Squared: 0.00241
Adjusted R−Squared: −0.000822857998501014
Root MSE: 708.609712488194
AIC: 9905.98185584123



minutes_played against age
minutes_played
0
1000
2000
3000
4000
20
25
30
35
age



minutes_played against injured
minutes_played
0
1000
2000
3000
4000
0.0
0.2
0.4
0.6
0.8
1.0
injured



Residuals vs Fitted
Residuals
1000
0
−1000
−2000
2500
2550
2600
2650
Fitted values
lm(formula)
53
43
480


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
43
480
546


Scale−Location
43
480
546
√|Standardized residuals|
0.0
0.5
1.0
1.5
2500
2550
2600
2650
Fitted values
lm(formula)


Cook's distance
546
480
43
0.04
0.03
0.02
0.01
0.00
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
43
480
546
0.000
0.005
0.010
0.015
−4
−3
−2
−1
0
1
2


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
546
480
43
Cook's distance
0.00
0.01
0.02
0.03
0.04
0
0.5
1
1.5
2
2.5
0
0.005
0.01
0.015
Leverage h_{ii}
lm(formula)