


minutes_played ~ age+prev_injuries
2616.62 + (−3.021)*age + (−0.697)*prev_injuries


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 0.08405 on 2 and 617 Degrees of Freedom
F p−value: 0.9194
R−Squared: 0.00027
Adjusted R−Squared: −0.00296824048591615
Root MSE: 709.36880038492
AIC: 9907.30947692055



minutes_played against age
minutes_played
0
1000
2000
3000
4000
20
25
30
35
age



minutes_played against prev_injuries
minutes_played
0
1000
2000
3000
4000
0
1
2
3
4
prev_injuries



Residuals vs Fitted
Residuals
1000
0
−1000
−2000
53
480
43
2510
2520
2530
2540
2550
2560
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
43
480
53


Scale−Location
43
480
53
√|Standardized residuals|
0.0
0.5
1.0
1.5
2510
2520
2530
2540
2550
2560
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.000
0.005
0.010
0.015
0.020
97
546
613
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
613
97
546
2
1
0
−1
−2
−3
−4
0.000
0.005
0.010
0.015
0.020


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
2
613
546
97
Cook's distance
0.000
0.005
0.010
0.015
0.020
0
0.5
1
1.5
0
0.005
0.01
0.015
0.02
Leverage h_{ii}
lm(formula)