

minutes_played ~ age+training_load+bmi+injured


2373.381 + (−6.016)*age + (149.337)*training_load + (−7.095)*bmi + (34.759)*injured


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05, High R and Adj R,
Low AIC


F−statistic: 7.53055 on 4 and 615 Degrees of Freedom
F p−value: 1e−05
R−Squared: 0.04669
Adjusted R−Squared: 0.0404918744087182
Root MSE: 692.704162119955
AIC: 9881.83151798412



minutes_played against age
minutes_played
0
1000
2000
3000
4000
20
25
30
35
age



minutes_played against training_load
minutes_played
0
1000
2000
3000
4000
1
2
3
4
5
training_load



minutes_played against bmi
minutes_played
0
1000
2000
3000
4000
20
22
24
26
28
bmi



minutes_played against injured
minutes_played
0
1000
2000
3000
4000
0.0
0.2
0.4
0.6
0.8
1.0
injured



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
2000
1000
0
−1000
−2000
2200
2300
2400
2500
2600
2700
2800
43
53
480


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
480
43
53
−3
−2
−1
0
1
2
3


Scale−Location
43
480
53
√|Standardized residuals|
0.0
0.5
1.0
1.5
2200
2300
2400
2500
2600
2700
2800
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0.035
43
480
546
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
3
2
1
0
−1
−2
−3
0.000
0.005
0.010
0.015
0.020
0.025
Leverage
lm(formula)
Cook's distance
480
43
546


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
1
0.5
0
0.5
1
1.5
2
2.5
480
43
546
Cook's distance
Leverage h_{ii}
lm(formula)
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0.035
0
0.005
0.01
0.015
0.02
0.025