

minutes_played ~ age+training_load+prev_injuries+bmi+injured


371 + ( −6.107)*age + ( 149.785)*training_load + ( −12.868)*prev_injuries + ( −6.972)*bmi + ( 43.287)*i


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 6.06383 on 5 and 614 Degrees of Freedom
F p−value: 2e−05
R−Squared: 0.04706
Adjusted R−Squared: 0.0392959871133971
Root MSE: 692.571950956645
AIC: 9883.59482604746



minutes_played against age
minutes_played
0
1000
2000
3000
4000
20
25
30
35
age



minutes_played against training_load
minutes_played
0
1000
2000
3000
4000
1
2
3
4
5
training_load



minutes_played against prev_injuries
minutes_played
0
1000
2000
3000
4000
0
1
2
3
4
prev_injuries



minutes_played against bmi
minutes_played
0
1000
2000
3000
4000
20
22
24
26
28
bmi



minutes_played against injured
minutes_played
0
1000
2000
3000
4000
0.0
0.2
0.4
0.6
0.8
1.0
injured



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
2000
1000
0
−1000
−2000
2200
2300
2400
2500
2600
2700
2800
2900
43
53
480


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
480
43
53
−3
−2
−1
0
1
2
3


Scale−Location
43
480
53
√|Standardized residuals|
0.0
0.5
1.0
1.5
2200
2300
2400
2500
2600
2700
2800
2900
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
43
480
546
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
3
2
1
0
−1
−2
−3
546
480
43
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
43
546
480
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0
0.5
1
1.5
2
0
0.005
0.01
0.015
0.02
0.025
Leverage $h_{ii}$
lm(formula)