


minutes_played ~ age+training_load
2192.562 + (−5.476)*age + (151.475)*training_load


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 14.91045 on 2 and 617 Degrees of Freedom
F p−value: 0
R−Squared: 0.0461
Adjusted R−Squared: 0.0430117480909837
Root MSE: 692.917924649184
AIC: 9878.21411225838



minutes_played against age
minutes_played
0
1000
2000
3000
4000
20
25
30
35
age



minutes_played against training_load
minutes_played
0
1000
2000
3000
4000
1
2
3
4
5
training_load



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
2000
1000
0
−1000
−2000
2200
2300
2400
2500
2600
2700
2800
43
53
480


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
43
480
53


Scale−Location
43
480
53
√|Standardized residuals|
0.0
0.5
1.0
1.5
2200
2300
2400
2500
2600
2700
2800
Fitted values
lm(formula)


Cook's distance
43
466
546
Cook's distance
0.00
0.01
0.02
0.03
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
3
2
1
0
−1
−2
−3
Cook's distance
0.000
0.005
0.010
0.015
Leverage
lm(formula)
546
466
43


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
43
546
466
Cook's distance
0.00
0.01
0.02
0.03
0
0.5
1
1.5
2
0
0.005
0.01
0.015
Leverage h_{ii}
lm(formula)