


minutes_played ~ bmi+injured
2775.48 + (−10.613)*bmi + (77.757)*injured


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 0.73655 on 2 and 617 Degrees of Freedom
F p−value: 0.47918
R−Squared: 0.00238
Adjusted R−Squared: −0.000851938006332764
Root MSE: 708.620007130318
AIC: 9905.99987036076



minutes_played against bmi
minutes_played
0
1000
2000
3000
4000
20
22
24
26
28
bmi



minutes_played against injured
minutes_played
0
1000
2000
3000
4000
0.0
0.2
0.4
0.6
0.8
1.0
injured



Residuals vs Fitted
Residuals
1000
0
−1000
−2000
43
466
480
2500
2550
2600
2650
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
43
480
466


Scale−Location
43
466
480
√|Standardized residuals|
0.0
0.5
1.0
1.5
2500
2550
2600
2650
Fitted values
lm(formula)


Cook's distance
43
480
546
0.00
0.01
0.02
0.03
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
546
480
43
0.000
0.005
0.010
0.015
0.020
−4
−3
−2
−1
0
1
2


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
43
480
546
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0.035
0
0.5
1
1.5
2
0
0.005
0.01
0.015
0.02
Leverage h_{ii}
lm(formula)