


minutes_played ~ injured
2525.61 + (79.016)*injured


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 1.16032 on 1 and 618 Degrees of Freedom
F p−value: 0.28182
R−Squared: 0.00187
Adjusted R−Squared: 0.000258934920428278
Root MSE: 708.800334814139
AIC: 9904.31538203013



minutes_played against injured
minutes_played
0
1000
2000
3000
4000
0.0
0.2
0.4
0.6
0.8
1.0
injured



Residuals vs Fitted
Residuals
1000
0
−1000
−2000
53
43
480
2540
2560
2580
2600
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
43
480
53


Scale−Location
43
53
480
√|Standardized residuals|
0.0
0.5
1.0
1.5
2540
2560
2580
2600
Fitted values
lm(formula)


Cook's distance
0.04
0.03
0.02
0.01
0.00
Cook's distance
480
546
97
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
97
546
480
0.000
0.002
0.004
0.006
0.008
2
1
0
−1
−2
−3
−4


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
480
546
97
Cook's distance
0.00
0.01
0.02
0.03
2.5
2
1.5
1
0.5
0
0.001
0.002
0.003
0.004
0.005
0.006
0.007
0.008
0.009
Leverage h_{ii}
lm(formula)