

minutes_played ~ prev_injuries+bmi+injured


2779.932 + (−7.827)*prev_injuries + (−10.543)*bmi + (82.957)*injured


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 0.51806 on 3 and 616 Degrees of Freedom
F p−value: 0.66998
R−Squared: 0.00252
Adjusted R−Squared: −0.00234119982493652
Root MSE: 708.572115886656
AIC: 9907.9160635978



minutes_played against prev_injuries
minutes_played
0
1000
2000
3000
4000
0
1
2
3
4
prev_injuries



minutes_played against bmi
minutes_played
0
1000
2000
3000
4000
20
22
24
26
28
bmi



minutes_played against injured
minutes_played
0
1000
2000
3000
4000
0.0
0.2
0.4
0.6
0.8
1.0
injured



Residuals vs Fitted
Residuals
1000
0
−1000
−2000
2500
2550
2600
2650
Fitted values
lm(formula)
43
466
480


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
43
480
466


Scale−Location
43
466
480
√|Standardized residuals|
0.0
0.5
1.0
1.5
2500
2550
2600
2650
Fitted values
lm(formula)


Cook's distance
43
220
480
0.000
0.005
0.010
0.015
0.020
0.025
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
220
480
43
2
1
0
−1
−2
−3
−4
0.000
0.005
0.010
0.015
0.020
0.025


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
43
480
220
Cook's distance
Leverage h_{ii}
lm(formula)
0.000
0.005
0.010
0.015
0.020
0.025
0
0.5
1
1.5
2