


minutes_played ~ prev_injuries+injured
2531.868 + (−8.021)*prev_injuries + (84.336)*injured


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 0.62308 on 2 and 617 Degrees of Freedom
F p−value: 0.53663
R−Squared: 0.00202
Adjusted R−Squared: −0.00121930733646791
Root MSE: 708.750047030982
AIC: 9906.22740370931



minutes_played against prev_injuries
minutes_played
0
1000
2000
3000
4000
0
1
2
3
4
prev_injuries



minutes_played against injured
minutes_played
0
1000
2000
3000
4000
0.0
0.2
0.4
0.6
0.8
1.0
injured



Residuals vs Fitted
Residuals
1000
0
−1000
−2000
53
43
480
2500
2520
2540
2560
2580
2600
2620
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
43
480
53


Scale−Location
43
53
480
√|Standardized residuals|
0.0
0.5
1.0
1.5
2500
2520
2540
2560
2580
2600
2620
Fitted values
lm(formula)


Cook's distance
97
480
613
0.000
0.005
0.010
0.015
0.020
0.025
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
613
480
97
0.000
0.005
0.010
0.015
−4
−3
−2
−1
0
1
2


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
480
613
97
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0
0.005
0.01
0.015
2
1.5
1
0.5
0
Leverage h_{ii}
lm(formula)