


minutes_played ~ prev_injuries
2541.012 + (−0.825)*prev_injuries


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 0.00098 on 1 and 618 Degrees of Freedom
F p−value: 0.97501
R−Squared: 0
Adjusted R−Squared: −0.00161653078053292
Root MSE: 709.46486081357
AIC: 9905.47738234169



minutes_played against prev_injuries
minutes_played
0
1000
2000
3000
4000
0
1
2
3
4
prev_injuries



Residuals vs Fitted
Residuals
1000
0
−1000
−2000
480
53
43
2538.0
2538.5
2539.0
2539.5
2540.0
2540.5
2541.0
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
43
480
53


Scale−Location
43
480
53
√|Standardized residuals|
0.0
0.5
1.0
1.5
2538.0
2538.5
2539.0
2539.5
2540.0
2540.5
2541.0
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
97
220
613
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
220
613
97
2
1
0
−1
−2
−3
−4
0.000
0.005
0.010
0.015


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
613
97
220
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0
0.002
0.004
0.006
0.008
0.01
0.012
0.014
Leverage h_{ii}
lm(formula)
2
1.5
1
0.5
0