

minutes_played ~ training_load+prev_injuries+bmi


2232.152 + (150.361)*training_load + (−9.467)*prev_injuries + (−7.004)*bmi


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05, High R and Adj R,
Low AIC


F−statistic: 9.82185 on 3 and 616 Degrees of Freedom
F p−value: 0
R−Squared: 0.04565
Adjusted R−Squared: 0.0410022482497993
Root MSE: 693.082704433482
AIC: 9880.50895618443



minutes_played against training_load
minutes_played
0
1000
2000
3000
4000
1
2
3
4
5
training_load



minutes_played against prev_injuries
minutes_played
0
1000
2000
3000
4000
0
1
2
3
4
prev_injuries



minutes_played against bmi
minutes_played
0
1000
2000
3000
4000
20
22
24
26
28
bmi



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
2000
1000
0
−1000
−2000
2200
2300
2400
2500
2600
2700
2800
43
53
480


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
43
480
53


Scale−Location
43
480
53
√|Standardized residuals|
0.0
0.5
1.0
1.5
2200
2300
2400
2500
2600
2700
2800
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.00
0.01
0.02
0.03
0.04
43
220
613
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
613
220
43
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
2
1
0
−1
−2
−3


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
43
220
613
2
1.5
1
0.5
0
Cook's distance
0.00
0.01
0.02
0.03
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)