opponent_rank ~ goals_conceded
15.418 + (−3.348)*goals_conceded

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC

F−statistic: 7.22202 on 1 and 44 Degrees of Freedom
F p−value: 0.01013
R−Squared: 0.14099
Adjusted R−Squared: 0.121471516433526
Root MSE: 6.51712173004654
AIC: 308.990165048936


opponent_rank against goals_conceded
opponent_rank
goals_conceded
5
10
15
20
0.0
0.5
1.0
1.5
2.0
2.5
3.0


Residuals vs Fitted
Residuals
Fitted values
lm(formula)
31
1
3
10
5
0
−5
−10
6
8
10
12
14


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−2
−1
0
1
2
3
1
31


Scale−Location
√|Standardized residuals|
0.0
0.2
0.4
0.6
0.8
1.0
1.2
6
8
10
12
14
Fitted values
lm(formula)
31
1
3


Cook's distance
Cook's distance
0.00
0.02
0.04
0.06
0.08
0.10
3
11
44
0
10
20
30
40
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
0.5
0.5
44
11
3
2
1
0
−1
−2
0.00
0.05
0.10
0.15


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
Cook's distance
Leverage h_{ii}
lm(formula)
0.00
0.02
0.04
0.06
0.08
0.10
0
0.05
0.1
0.15
2
1.5
1
0.5
0
44
3
11