opponent_rank ~ goals_scored+goals_conceded
13.172 + (1.448)*goals_scored + (−2.915)*goals_conceded

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC

F−statistic: 6.11051 on 2 and 43 Degrees of Freedom
F p−value: 0.00462
R−Squared: 0.22131
Adjusted R−Squared: 0.18509306474976
Root MSE: 6.20497171548574
AIC: 306.474624202783


opponent_rank against goals_scored
opponent_rank
goals_scored
5
10
15
20
0
2
4
6
8


opponent_rank against goals_conceded
opponent_rank
goals_conceded
5
10
15
20
0.0
0.5
1.0
1.5
2.0
2.5
3.0


Residuals vs Fitted
Residuals
Fitted values
lm(formula)
44
46
1
10
5
0
−5
−10
−15
10
15
20
25


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−2
−1
0
1
2
1
44
46


Scale−Location
√|Standardized residuals|
0.0
0.2
0.4
0.6
0.8
1.0
1.2
1.4
44
46
1
10
15
20
25
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.00
0.02
0.04
0.06
0.08
0.10
3
11
44
0
10
20
30
40
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
1
0.5
0.5
1
44
3
11
2
1
0
−1
−2
0.0
0.1
0.2
0.3
0.4
0.5


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
2
1.5
1
0.5
44
11
3
Cook's distance
0.00
0.02
0.04
0.06
0.08
0.10
0
0.1
0.2
0.3
0.4
0.5
Leverage h_{ii}
lm(formula)