opponent_rank ~ goals_scored
10.401 + (1.726)*goals_scored

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC

F−statistic: 5.85895 on 1 and 44 Degrees of Freedom
F p−value: 0.01969
R−Squared: 0.11751
Adjusted R−Squared: 0.0974539588578468
Root MSE: 6.60560488077487
AIC: 310.23084825131


opponent_rank against goals_scored
opponent_rank
goals_scored
5
10
15
20
0
2
4
6
8


Residuals vs Fitted
Residuals
Fitted values
lm(formula)
46
29
1
10
5
0
−5
−10
−15
10
12
14
16
18
20
22
24


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
46
29
1
2
1
0
−1
−2
−2
−1
0
1
2


Scale−Location
√|Standardized residuals|
0.0
0.2
0.4
0.6
0.8
1.0
1.2
1.4
46
29
1
10
12
14
16
18
20
22
24
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.00
0.01
0.02
0.03
0.04
0.05
0.06
0.07
1
31
46
0
10
20
30
40
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
46
31
1
0.5
1
0.5
1
2
1
0
−1
−2
0.0
0.1
0.2
0.3
0.4
0.5


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
2
1.5
1
0.5
46
1
31
0
Cook's distance
0.00
0.01
0.02
0.03
0.04
0.05
0.06
0
0.1
0.2
0.3
0.4
0.5
Leverage h_{ii}
lm(formula)