prev_injuries ~ age+bmi+injured

0.741 + (−0.007)*age + (0.009)*bmi + (0.674)*injured

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC

F−statistic: 12.48137 on 3 and 616 Degrees of Freedom


F p−value: 0


R−Squared: 0.0573


Adjusted R−Squared: 0.0527116308246389


Root MSE: 1.05223732438811


AIC: 1832.62294734725



prev_injuries against age
prev_injuries
0
1
2
3
4
20
25
30
35
age



prev_injuries against bmi
prev_injuries
bmi
0
1
2
3
4
20
22
24
26
28



prev_injuries against injured
prev_injuries
injured
0
1
2
3
4
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
Residuals
3
2
1
0
−1
−2
0.8
1.0
1.2
1.4
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
534
−3
−2
−1
0
1
2
3


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.8
1.0
1.2
1.4
Fitted values
lm(formula)


Cook's distance
271
129
256
0.025
0.020
0.015
0.010
0.005
0.000
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
129
256
271
Cook's distance
3
2
1
0
−1
−2
0.000
0.005
0.010
0.015
0.020


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
271
1256
Cook's distance
0.025
0.020
0.015
0.010
0.005
0.000
2
1.5
1
0.5
0
0
0.005
0.01
0.015
0.02
Leverage h_{ii}
lm(formula)