


prev_injuries ~ age+injured
0.947 + (−0.007)*age + (0.673)*injured


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 18.70149 on 2 and 617 Degrees of Freedom
F p−value: 0
R−Squared: 0.05716
Adjusted R−Squared: 0.0540996708060193
Root MSE: 1.05231924881698
AIC: 1830.7194867258



prev_injuries against age
prev_injuries
0
1
2
3
4
20
25
30
35
age



prev_injuries against injured
prev_injuries
injured
0
1
2
3
4
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
Residuals
Fitted values
lm(formula)
3
2
1
0
−1
−2
0.8
1.0
1.2
1.4


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
129


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.8
1.0
1.2
1.4
Fitted values
lm(formula)


Cook's distance
256
439
181
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
439
256
181
Cook's distance
3
2
1
0
−1
−2
0.000
0.005
0.010
0.015


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
256
439
181
Cook's distance
Leverage h_{ii}
lm(formula)
0.000
0.005
0.010
0.015
0.020
0.025
0
0.005
0.01
0.015
0
0.5
1
1.5
2