


prev_injuries ~ age+minutes_played+bmi
0.731 + ( 0.003)*age + ( 0)*minutes_played + ( 0.004)*bmi


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 0.03525 on 3 and 616 Degrees of Freedom
F p−value: 0.99113
R−Squared: 0.00017
Adjusted R−Squared: −0.00469763339246421
Root MSE: 1.08365313038898
AIC: 1869.10272930177



prev_injuries against age
prev_injuries
0
1
2
3
4
20
25
30
35
age



prev_injuries against minutes_played
prev_injuries
minutes_played
0
1
2
3
4
0
1000
2000
3000
4000



prev_injuries against bmi
prev_injuries
bmi
0
1
2
3
4
20
22
24
26
28



Residuals vs Fitted
256
439
181
Residuals
3
2
1
0
−1
0.86
0.88
0.90
0.92
0.94
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
−3
−2
−1
0
1
2
3
−1
0
1
2
3
534
439
256


Scale−Location
256
439
534
√|Standardized residuals|
0.0
0.5
1.0
1.5
0.86
0.88
0.90
0.92
0.94
Fitted values
lm(formula)


Cook's distance
256
129
220
0.020
0.015
0.010
0.005
0.000
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
256
129
220
3
2
1
0
−1
0.000
0.005
0.010
0.015
0.020
0.025
0.030


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
2
1.5
1
0.5
0
256
129
220
Cook's distance
0.000
0.005
0.010
0.015
0.020
0
0.005
0.01
0.015
0.02
0.025
Leverage $h_{ii}$
lm(formula)